



COHESIVE HEALTHCARE MANAGEMENT & CONSULTING

Mangum Regional Medical Center

TITLE		POLICY	
Iodine Allergy and IV Contrast Pre-treatment Protocol		DRM-058	
MANUAL	EFFECTIVE DATE	REVIEW DATE	
Drug Room	10-1-2020	10-1-2020	
DEPARTMENT	REFERENCE		
Drug Room	<u>https://www.acr.org/</u>		

SCOPE

This protocol applies to adult patients that require treatment for an Iodine Allergy prior to IV contrast administration.

PROCEDURE

- Prednisone 50mg PO Once (Administer 13 hours prior to scan)
- Prednisone 50mg PO Once (Administer 7 hours prior to scan)
- Prednisone 50mg PO Once (Administer 1 hour prior to scan)
- Diphenhydramine 50mg PO Once (Administer 1 hour prior to scan)
- Diphenhydramine 50mg IV Once (Administer 1 hour prior to scan)
- Hydrocortisone 200mg IV Once (Administer 1 hour prior to scan)
- Methylprednisolone 40mg IV Once (Administer 1 hour prior to scan)

REFERENCES

<https://www.acr.org/>

ATTACHMENTS

None.

REVISIONS/UPDATES

Date	Brief Description of Revision/Change