

COHESIVE HEALTHCARE MANAGEMENT & CONSULTING HOSPITAL NAME

Casirivimab/Imdevimab (Combination Therapy)							
Emergency Use Authorization (EUA) Standing Orders							
All items with an autocheck "\" are automatically initiated							
Name:	Date:		Time:				
Date of Birth:							
Allergies:		Code Status:		Wt:			
		DNR					
Initial halamin the haw her as ab	DNI	(:m:4:al aaa	h :4 area				
Initial below in the box by each item:	I certify the patient/legal represen below):	tauve was	(шппат еас	ii iteiii			
	rnatives to Casirivimab/Imdevimab.						
	Parents, and Caregivers" prior to adm	inistration.					
The patient meets the appropriate c	riteria for administration (check each	item as app	olicable):				
$\Box \ge 12$ years of age	$\Box \ge 40 \text{ kg (weight)}$			moderate			
			COVID-1	9 disease			
☐ At high risk for progressing to severe COVID-19 and/or hospitalization.							
□ NOT hospitalized due to COVID							
□ <i>DO NOT</i> require oxygen therapy due to COVID-19, or □ <i>DO NOT</i> require an increase in baseline oxygen flow rate due to COVID-19 in those on chronic oxygen							
		ID-19 in th	ose on chro	onic oxygen			
therapy due to underlying non-COVID-19 related co-morbidity. Date of symptom onset: Date of positive test:							
Qualifying Reasons for Administration (Must choose at least one of the following):							
\Box BMI \geq 35 \Box Have chronic kidney disease \Box Diabetes							
☐ Immunosuppressive Disease	□ Currently receiving immunosuppr		□ Age ≥ 65 years				
treatment							
Are \geq 55 years of age AND have \square Cardiovascular disease, or \square Hypertension, or \square COPD/other chronic							
respiratory disease							
Are 12-17 years of age AND have (Check all that apply): \square BMI \ge 85 th percentile for their age and gender based							
on CDC growth charts, or □ Sickle Cell Disease, or □ Congenital or acquired heart disease, or							
□ Neurodevelopmental disorders, i.e., Cerebral Palsy, or □ Medical-related technological dependence, i.e.,							
tracheostomy, gastrostomy, or positive pressure ventilation (not related to COVID-19), or □ Asthma, reactive							
airway disease or other chronic respiratory disease that requires daily medication for control.							
ORDERS							
√ Casirivimab 600mg/Imdevimab 600mg IV infusion over 60 minutes as soon as possible after positive viral							
test for SARS-CoV-2 and within 10 days of symptom onset. Once the infusion is complete, flush the tubing with							
0.9% Sodium Chloride to ensure delivery of the dose.							
√ Administer infusion using 0.2 micron filter tubing.							
$\sqrt{\text{Obtain baseline VS (Temp, Pulse, Respiration, BP, O2 Sat)}}$ prior to infusion.							

Nurse Signature:					Ti	me:	Date:			□ TORB □ VORB	
Provid	ler Signature:				Ti	me:		Date:			
Do Not	Use Instead	Do Not	Use	Do Not	Use	Do Not	Use	Do Not	Use	Do Not	Use Instead
Use		Use	Instead	Use	Instead	Use	Instead	Use	Instead	Use	
U	Unit	1.0 mg	1 mg	QD	Daily	MS or	Morphine	сс	mL	SC,	Subcutaneous
						MSO4				SQ,	
										Sub q	
IU	International Unit	.X mg	0.X mg	QOD	Every	MgSO4	Magnesium	qhs	nightly		Discharge or
					Other Day		Sulfate			D/C	Discontinue

√ Monitor VS (Temp, Pulse, Respiration, BP, O2 Sat) every 30 minutes until one hour after infusion is										
complete. Notify Provider if patient exhibits any of the following signs or symptoms:										
Temp > 100.4°F	Chills	Nausea	Headache	Bronchospasm	Hypotension					
Angioedema	Throat Irritation	Rash/Urticaria	Pruritis	Myalgia	Dizziness					
□ Outpatient: Instruct patient to continue to self-isolate and use infection control measures according to CDC										
guidelines (i.e. wear a mask, social distance, avoid sharing personal items, clean & disinfect "high touch surfaces,"										
frequent hand hyg	frequent hand hygiene).									
		Allergic/Anaphy								
	on related to the info		the infusion. Initiat	e a Rapid Response	e or Code Blue as					
	tify the Provider im									
	ent of anaphylaxis:				r Trendelenburg					
position, administe	er supplemental oxy									
			sion (decreased cir							
□ Infuse 0.9% Nor			ain systolic BP > 90							
			y Distress (stridor							
	000 0.3mg IM or Su									
	and/or laryngeal ed	ema), hypotension,	and/or acute loss of	consciousness. Ma	ay repeat x1 in 10					
minutes if necessa		10								
	via nebulizer over	•	•	•						
☐ If wheezing pers	sists and BP is > 90r									
□ Acetaminophen	1000mg PO for hea		eadache, dizziness	, seizure)						
	et physician immedi									
	_ <u> </u>	<u> </u>	usea, emesis, diari	rhea)						
GI-(abdominal pain, nausea, emesis, diarrhea) □ Diphenhydramine 50mg IV or IM x1										
1		Skin-(rash, itchi	ing, welts, hives)							
□ Diphenhydramii	ne 50mg IV or IM fo									
□ Methylprednisol										
ADDITIONAL ORDERS										

Nurse Signature:					Ti	me:	Date:			□ TORB □ VORB	
Provid	ler Signature:				Ti	me:		Date:			
Do Not	Use Instead	Do Not	Use	Do Not	Use	Do Not	Use	Do Not	Use	Do Not	Use Instead
Use		Use	Instead	Use	Instead	Use	Instead	Use	Instead	Use	
U	Unit	1.0 mg	1 mg	QD	Daily	MS or	Morphine	сс	mL	SC,	Subcutaneous
						MSO4				SQ,	
										Sub q	
IU	International Unit	.X mg	0.X mg	QOD	Every	MgSO4	Magnesium	qhs	nightly		Discharge or
					Other Day		Sulfate			D/C	Discontinue