

Milkweeds (*Asclepias* spp.) are herbaceous perennial plants named for their milky sap. These plants occur in a wide range of habitats, including intact natural communities on roadsides and highly disturbed roadsides. As required host plants for monarch (*Danaus plexippus*) caterpillars, milkweeds play an essential role in the butterfly's life cycle (see reverse). Vegetation management that allows milkweeds to persist can support monarchs. This guide can help you recognize the most common native species found on roadsides in your region.



The most common milkweeds in roadsides in the Great Lakes Region (in alphabetical order):



PLANT: Upright, unbranched stems. **LEAVES:** Alternate; long and narrow; pointed at tips; with short hairs. **HABITAT:** Prairies, meadows, open woods, disturbed areas, railways. **SOILS:** Sandy, rocky to clay; dry-wet. **BLOOM:** Jun-Aug; green with purple or cream.



PLANT: One to many upright, branched stems; smooth or with short hairs. **LEAVES:** Opposite; lance-shaped or narrow; with few short hairs. **HABITAT:** Moist prairies, marshes, ditches, edges of ponds, lakes, streams. **SOILS:** Silty to loamy or clayey; moist–wet, tolerates some mesic. **BLOOM:** Jun–Aug; pink or light purple.



PLANT: One to many stout, upright, unbranched stems; usually with short dense hairs. **LEAVES:** Opposite; oval-shaped; hairy underneath. **HABITAT:** Prairies, old fields, railways, open woods, flood plains, disturbed areas. **SOILS:** Sandy to loamy, rocky or clayey; dry-wet. **BLOOM:** May-Aug; light purple or pink.



PLANT: One to many spreading to upright stems; with short hairs; lacks milky sap. **LEAVES:** Alternate; lance-shaped; hairy underneath. **HABITAT:** Prairies, old fields, open woods. **SOILS:** Sandy, loamy, rocky; dry-mesic. **BLOOM:** May-Aug; orange to red or yellow.

Most common milkweed species continued



PLANT: One to several upright, unbranched stems; with short hairs. LEAVES: Whorled; narrow to needle-like; smooth or with short hairs. HABITAT: Prairies, open woods, fields, flood plains, glades, railways, disturbed areas. SOILS: Sandy, rocky, clayey; dry-mesic. BLOOM: May-Sep; white to green.

Additional Resources:

- For more information on monarchs and roadsides, including monitoring, visit: <u>tinyurl.com/MJV-Monarchs-Roadsides</u>
- Mowing and Monarchs: tinyurl.com/MJV-MowingForMonarchs
- Serces Society for Invertebrate Conservation: xerces.org
- Monarch Joint Venture: monarchjointventure.org

THE MONARCH BUTTERFLY LIFE CYCLE 1 Egg 3-5 DAYS 2 Larva 10-14 DAYS Caterpillar grows by molting 5 INSTARS 3 Chrysalis 10-14 DAYS

Multiple generations of monarchs are produced over the spring and summer, with the fall generation migrating to overwintering sites. You can monitor monarchs or milkweeds; see Additional Resources above.

Less common roadside milkweeds:



Clasping milkweed (A. amplexicaulis)

PLANT: Upright, unbranched, stems; smooth; 3' max. LEAVES: Opposite; oval-shaped; wavy margins; base of leaves clasp stem. SOILS/HABITAT:



Sandy; dry; grasslands, savannas, woodland edges. **BLOOM:** Apr–Jun; pink or green with cream, sometimes purple.



Purple milkweed (A. purpurascens)

PLANT: Upright, unbranched, stout stems; smooth; 6' max. LEAVES: Opposite; oval-shaped; smooth above with fine hairs below. SOILS/



HABITAT: Loamy to clayey; mesic; prairies, meadows, woodland edges, savannas. **BLOOM:** May–Jul; dark rose to purple.



Prairie milkweed (A. sullivantii)

PLANT: Upright, unbranched, stout stems; smooth; 4' max. **LEAVES:** Opposite; lance- to oval-shaped; very smooth. **SOILS/HABITAT:** Sandy



clay or loamy; mesic-moist; prairies, wet meadows. **BLOOM:** Jun-Aug; pink with light green or purple.



Green comet milkweed (A. viridiflora)

PLANT: Multiple unbranched, spreading or upright stems; with short hairs; 3' max. LEAVES: Opposite; lance- or oval-shaped,



edges folded upward or wavy. SOILS/HABITAT: Sandy, rocky; dry; prairies, old fields, dunes, open woods. BLOOM: Jun-Jul; light to yellowish green.

Additional milkweeds in the Great Lakes region: Asclepias exaltata, A. lanuginosa, A. quadrifolia, A. ovalifolia, A. speciosa, A. stenophylla, A. variegata.

Maps & Distribution Data:

These profiles are derived from regional floras and field guides and Woodson's *The North American Species of* Asclepias (1954). Most common species are abundant across the states and are found in roadsides. Less common species might not occur in all states, have a limited distribution across a state, or may be less common in roadsides. Additional species may be uncommon in roadsides, have a small distribution in a state or region, or are uncommon or rare. The range maps indicate counties where species have been observed (but may be incomplete), and were created by USDA-NRCS using the latest data from the USDA's PLANTS database (https://plants.sc.egov.usda.gov).

ACKNOWLEDGMENTS: Written by Jennifer Hopwood, Stephanie Frischie, and Karin Jokela (Xerces Society), and Alison Cariveau (Monarch Joint Venture). Reviewed by Dan Zay (Michigan NRCS). Design, header, and monarch life cycle by Sara Morris (Xerces Society). This work was conducted in the National Cooperative Highway Research Program, which is administered by the Transportation Research Board of the National Academies of Sciences, Engineering, and Medicine.

PHOTO CREDITS: Jim Fowler (*A. syriaca*); Krista Lundgren, USFWS / flickr (*A. viridiflora*); Joshua Mayer / flickr (*A. hintellà*); Jerry Oldenettel / flickr (*A. verticillatà*); Tom Potterfield / flickr (*A. incamata, A. purpurascens*); Paul Rothrock / SEINet (*A. amplexicaulis*); Scott Seigfreid (*A. tuberosa*), Xerces Society / Ray Moranz (*A. sullivantil*). Photographs remain under the copyright of the photographer. © 2019 by The Xerces Society for Invertebrate Conservation. Xerces® is a trademark registered in the U.S. Patent and Trademark Office.

19-021 O1